

IOBC/wprs Bulletin Vol. 28(7) 2005

Working Groups „Integrated fruit protection in fruit crops” and “Use of pheromones and other semiochemicals in integrated control”. Proceedings of the 6th International Conference on Integrated Fruit Production. Baselga di Piné (Italy), September 26-30, 2004. Edited by: Jerry Cross & Claudio Ioriatti. ISBN 92-9067-179-9 [xxviii + 485 pp.]

IFP

| | |
|---|----|
| Strategies to control apple proliferation disease in Trentino and in Germany <i>Wolfgang Jarausch, Annamaria Ciccotti, Luisa Mattedi, Elisabetta Vindimian</i> | 3 |
| The third edition of the IOBC Basic Documents on Integrated Production <i>Ernst F. Boller, Jesús Avilla, Erich Jörg, Carlo Malavolta, Frank G. Wijnands, Peter Esbjerg</i> | 9 |
| Controlled Integrated Production of Fruit – A Comparison of Production Guidelines and Checking Procedures in Europe <i>Bernhard Sessler, Graciela Wiegand, Tilman Becke</i> | 13 |
| A history and achievements of IOBC subgroup - Soft Fruits <i>Dariusz Gajek, Jerry Cross, Erich Jörg</i> | 17 |
| Development of an integrated pest and disease management system for apples to produce fruit free from pesticide residues <i>Angela Berrie, Jerry Cross</i> | 22 |
| Interaction between fruit tree canker and apple scab control in integrated apple orchards <i>Bart Heijne, Peter Frans de Jong, Imre Holb</i> | 33 |
| Preventive apple scab infection warnings: optimization of leaf wetness models and evaluation of regional weather forecast data. <i>Stijn Van Laer, Piet Creemers</i> | 37 |
| Assessing the risk of reduced-risk apple pest control programs in New York State <i>Arthur Agnello, Jan Nyrop, Harvey Reissig, Richard Straub</i> | 43 |
| Development of a new paradigm for management of internal lepidoptera in western New York apple orchards <i>Harvey Reissig, Art Agnello, Jan Nyrop</i> | 49 |
| Ecologically determined differences in the insect-host plant relations between various populations of codling moth <i>Asya Ter-Hovhannesyan</i> | 55 |
| Spatial analysis of pheromone trap catches of <i>Cydia funebrana</i> , <i>Cydia pomonella</i> , <i>Cydia molesta</i> and <i>Anarsia lineatella</i> : contribution to the IPM in fruit crops <i>Andrea Sciarretta, Mossbah Alarram, Pasquale Trematerra</i> | 61 |
| The potential of integrating the Sterile Insect Technique as an environmentally friendly method for area-wide management of the codling moth (<i>Cydia pomonella</i>) <i>Marc Vreysen and Jorge Hendrichs</i> | 65 |
| Components of an ecologically and economically sustainable orchard <i>Mark W. Brown and Clarissa R. Mathews</i> | 73 |
| Role of leaf miners parasitoids in an IPM sour cherry plantations <i>Klára Balázs, Krisztina Mihályi, Viktor Markó, Csaba Thuróczy</i> | 77 |
| Reduction of insecticide spraying by using alternative methods in commercial apple orchards <i>J.L. Batllori, M. Vilajeliu, P. Vilardell, A. Creixell, M. Carbó, N. Garcia, G. Esteba, F. Raset, F. Vayreda, M. Giné, D. Curós, J. Cornella</i> | 83 |
| The Tortricidae population in Croatian apple orchards and a possibility for mating disruption techniques <i>Božena Barić, Ivan Ciglar</i> | 89 |
| Development of IPM programs in apple orchards by autosterilization of codling moth wild populations | |

| | |
|--|-----|
| Asya Ter-Hovhannesyanyan, Anzhela Azizyan | 93 |
| Pilot project for sustained management of codling moth, <i>Cydia pomonella</i> L., with minimal SIT, in British Columbia, 2001– 2003 Howard Thistlewood, Gary Judd, Markus Clodius | 99 |
| The egg stage as the target for the control of the Oriental Fruit Moth (<i>Cydia molesta</i> (Busck); Lepidoptera: Tortricidae) Emanuele Mazzoni, Fabio Molinari | 105 |
| Seasonal fluctuation of Oriental Fruit Moth (<i>Grapholita molesta</i> (Busck)) with sexual pheromone trap in peach orchards in Bento Gonçalves, RS, Brazil Cristiano João Arioli, Marcos Botton, Geraldo Andrade Carvalho | 111 |
| A forecasting model for the peach twig borer, <i>Anarsia lineatella</i> (Zeller) F. Molinari, R. Tiso, A. Butturini, G. Ceredi, P. Sambado, E. Rossi | 115 |
| Occurrence of visual symptoms of pest and disease in 15 sweet cherry varieties under organic cultivation Alessandro Roversi, Antonio Monteforte | 119 |
| Preliminary study of <i>Forficula</i> sp. (Dermaptera: Forficulidae) as predator of <i>Cacopsylla pyri</i> (Homoptera: Psyllidae), under field conditions in Lleida (Spain). Jauset, A. M., Artigues, M., Avilla J., Sarasúa, M. J. | 125 |
| Sustainable control of the rosy apple aphid <i>Dysaphis plantaginea</i> Marcos Miñarro, Enrique Dapena | 129 |
| Preliminary investigation on the natural enemies of the peach twig borer, <i>Anarsia lineatella</i> in northern Italy Fabio Molinari, Elisabetta Chiappini, Paolo Sambado | 135 |
| Occurrence of the anthocorids <i>Anthocoris nemorum</i> and <i>A. nemoralis</i> in apple and pear in Denmark Lene Sigsgaard | 139 |
| Fly-free cherries: utopia or realistic hope? Kirsten Köppler, Arne Peters, Volker Storch, Heidrun Vogt | 143 |
| Effects of plant protection products on <i>Kampimodromus aberrans</i> (Oudemans): the dietary effect of airborne pollen Mario Baldessari, Gino Angeli, Vincenzo Girolami | 151 |
| Bioassay methodology and resistance to insecticides of pear psylla (<i>Cacopsylla pyri</i> L.) (Homoptera: Psyllidae) in Lleida, Spain Xavier Miarnau, María José Sarasúa, Jesus Avilla, Miquel Artigues | 157 |
| Organic cider-apple production in Asturias (NW Spain) Enrique Dapena, Marcos Miñarro, María Dolores Blázquez | 161 |
| Hazelnut fruit quality as influenced by the cultivation system adopted Alessandro Roversi, Claudio Sonnati | 167 |
| Digital analysis of injuries caused by raspberry spider mite <i>Neotetranychus rubi</i> (Träg.) and two-spotted spider mite <i>Tetranychus urticae</i> (Koch.) on raspberry leaves Dariusz Gajek, Wojciech Warabieda, Remigiusz W. Olszak | 173 |
| Temperature model of the flight activity of <i>Adoxophyes orana</i> (Lep.: Tortricidae) for timing of control Jitka Stará, František Kocourek | 177 |
| Monitoring of San Jose scale (<i>Quadraspidiotus perniciosus</i> Comst.) by pheromone traps and timing of control on crawlers Kocourek František, Stará Jitka | 183 |
| Secondary dissemination of <i>Bacillus subtilis</i> (BD170 - BioPro) against <i>Erwinia amylovora</i> on apple flowers by means of pollinators, <i>Apis mellifera</i> and <i>Osmia cornuta</i> Maccagnani Bettina, Betti Federica, Fanti Marco, Davide Gobbin, Pertot Ilaria, Angeli Gino, Gessler Cesare | 189 |
| Biological efficacy of kaolin against the pear sucker <i>Psylla pyri</i> in winter and summer applications Bruno Gobin, Dany Bylemans, Gertie Peusens | 193 |
| Effects of some botanical pesticides on pests and beneficial arthropods Teodoro Moleas, Serafina Di Gioia, Rocco Addante | 199 |

| | |
|--|-----|
| Monitoring of the spread of AP subtypes in Trentino <i>Christian Cainelli, Stella Grando</i> | 205 |
| Resistant rootstocks as a strategy to control AP disease <i>Claudia Bisognin, Stella Grando</i> | 211 |
| The use of a 'visual+temperature' method in timing of cherry fruit fly (<i>Rhagoletis cerasi</i> L.) control <i>Vladan Falta, Václav Kneifl, Jaroslava Křourková</i> | 217 |
| Control of phytophagous mites on strawberry in Europe by predatory phytoseiid mites or heat treatment <i>Jean Fitzgerald, Dany Bylemans, Gertie Peusens, Nina Trandem, Tuomo Tuovinen</i> | 221 |
| Prey preferences of <i>Anthocoris nemoralis</i> and <i>A. nemorum</i> (Heteroptera: Anthocoridae) and their predation behaviour towards Pear psyllid, <i>Cacopsylla pyri</i> <i>Lene Sigsgaard</i> | 225 |
| The effect of different growing methods on the incidence of cane pests in Hungarian raspberry plantations <i>Gábor Véték, Béla Péntzes</i> | 229 |
| Use of a New Generation of Horticultural Oils for Mite Management in Fruit Orchards. <i>Greg Krawczyk, Larry A. Hull</i> | 233 |
| The judgement of the population regulating effect of <i>Zetzellia mali</i> (Stigmaeidae) <i>Gábor Jenser</i> | 239 |
| Effects of modern bait formulated pesticides on larvae and adults of <i>Chrysoperla carnea</i> under extended-laboratory conditions <i>G. Contreras, P. Medina, A. Adán, A.N. Zapata, E. Viñuela</i> | 245 |
| Reduced application rates of imidacloprid on apple: effect on leafhoppers, aphids and aphid predators <i>Richard W. Straub, Peter J. Jentsch</i> | 251 |
| Trials on the efficacy of natural products against Oriental Fruit Moth in organic peach orchard. <i>Fabio Molinari, Emanuele Mazzoni, Paolo Sambado, Piero Cravedi</i> | 255 |
| Control of Codling moth <i>Cydia pomonella</i> L. using insecticides: relationship between susceptibility and efficacy in the field. <i>Claudio Ioriatti, Pierre-Joseph Charmillot, Flavia Forno, Luisa Mattedi, Denis Pasquier, Claudio Rizzi</i> | 259 |
| Detection of codling moth (<i>Cydia pomonella</i>) resistance by topical application of insecticides and validation on a laboratory resistant strain by dipping of apples and incorporating products into artificial diets <i>P. J. Charmillot, D. Pasquier, F. Briand</i> | 265 |
| Biological control of <i>Metcalfa pruinosa</i> with <i>Neodrynus typhlocybae</i> : on establishment and diffusion of the parasitoid in Trentino Alto Adige (Italy) <i>Gino Angeli, Romano Maines, Marco Fantì, Diego Forti, Monica Sofia, Mario Baldessari, Cristina Tomasi, Oliviero Sandri, Lodovico Delaiti, Claudio Ioriatti, Vincenzo Girolami</i> | 271 |
| PHEROMONES | |
| Constant Monitoring Enhances of Pheromones in IFP <i>Walter Waldner</i> | 277 |
| Mating disruption to control grapevine moth, <i>Lobesia botrana</i> (Den. and Schiff.) in Porto Wine Region: a three-year study <i>C. Carlos, J. Costa, C. Gaspar, J. Domingos, F. Alves, L. Torres</i> | 283 |
| Codling moth management: from I.H.E.L.P. to M.A.P.S. to A.K.I.S.S. <i>Alan Knight</i> | 289 |
| The parasitoids of the European grapevine moth (<i>Lobesia botrana</i> Den.-Schiff.) and predators in the mating disruption-treated vineyards in Turkey <i>Turkan Koclu, Ferhunde Ozlem Altindisli, Fatma Ozsemerci</i> | 293 |

| | |
|---|-----|
| Pheromone release by individual females of <i>Lobesia botrana</i> (Den. et Schiff.) (Lepidoptera Tortricidae) and their competition with pheromone dispensers <i>Gianfranco Anfora, Marco Tasin, Silvia Carlin, Giacinto S. Germinara, Claudio Ioriatti, Antonio De Cristofaro</i> | 299 |
| New biodegradable controlled-release pheromone dispenser for mating disruption of European grapevine moth <i>Lobesia botrana</i> Denis and Schiffermüller (Lepidoptera: Tortricidae) <i>Beatriu Femenia-Ferrer, Pilar Moya, Vicente Navarro-Llopis, Jaime Primo</i> | 305 |
| Mechanisms of the Exosect Auto-Confusion Technique <i>Philip Howse, Ken MacDonald</i> | 309 |
| Attraction of four tortricid moth species to high dosage pheromone rope dispensers: Observations implicating false plume following as an important factor in mating disruption <i>Lukasz L. Stelinski, Larry J. Gut, David Epstein, James R. Miller</i> | 313 |
| Semiochemical driven autodissemination of <i>Cydia pomonella</i> and <i>Adoxophyes orana</i> baculoviruses <i>J.V. Cross, D. Winstanley, N. Naish, S. Hilton, G. Keane, R. van Wezel, D. Gajek</i> | 319 |
| A reliable field test for the efficiency of mating disruption techniques <i>Eric Doye, Uwe T. Koch</i> | 325 |
| New developments in EAG techniques for field pheromone and plant odour measurements <i>Martin Gabriel, Jörg Zastra, Uwe T. Koch</i> | 329 |
| Survey of pheromone emission from different kinds of dispensers used for mating disruption in orchards and vineyards <i>D. Pasquier, P. J. Charmillot</i> | 335 |
| Kairomone-augmented mating disruption control for codling moth in Californian walnuts and apples <i>Douglas Light, Alan Knight</i> | 341 |
| Season-long control of Oriental fruit moth by mating disruption in apples <i>Orkun Kovanci, James Walgenbach, George Kennedy</i> | 345 |
| Plant odours influence the host finding behaviour of apple psyllids (<i>Cacopsylla picta</i> ; <i>C. melanoneura</i>) <i>Jürgen Gross, Negash Mekonen</i> | 351 |
| Tortricid species caught by the Codling moth kairomone ethyl (2E,4Z)-2,4-decadienoate: monitoring trials and electrophysiological responses <i>Silvia Schmidt, Gianfranco Anfora, Antonio De Cristofaro, Claudio Ioriatti</i> | 357 |
| Potential for disruption of mate-seeking <i>Sesamia nonagrioides</i> (Lef.) males by (Z)-9-tetradecenyl acetate <i>G.S. Germinara, G. Rotundo, A. de Cristofaro, A. Elgargoti</i> | 363 |
| Control of the Codling moth, <i>Cydia pomonella</i> (L.) (Lepidoptera Tortricidae), by disorientation <i>Gino Angeli, Gianfranco Anfora, Mario Baldessari, Antonio De Cristofaro, Giacinto S. Germinara, Marco Tasin, Silvia Vitagliano, Claudio Ioriatti</i> | 371 |
| Impact of Increased point source densities on communication disruption of tortricid moth pests in Michigan tree fruit <i>David Epstein, Larry Gut, Lukasz Stelinski, James Miller</i> | 379 |
| Mating disruption for the control of European grapevine moth <i>Lobesia botrana</i> (Den. et Schiff.) in a plastic film greenhouse table grape vineyard <i>F. Savino, A. Iodice, V. Veronelli, K. Ogawa, T. Kobayashi</i> | 385 |
| Preliminary investigation on the mating behaviour of the peach twig borer, <i>Anarsia lineatella</i> . <i>Davide Natale, Fabio Molinari, Paolo Sambado, Piero Cravedi</i> | 391 |
| The use of the aphid sex pheromone and plant volatiles to enhance control of <i>Dysaphis plantaginea</i> in apple | |

| | |
|---|-----|
| <i>Jean Fitzgerald, Tom Pope, Mike Solomon, Guy Poppy, Alex Stewart Jones, Lester Wadhams</i> | 395 |
| Impact of the kairomone ethyl (2E, 4Z)-2,4-decadienoate (DA 2313) on the oviposition behaviour of <i>Cydia pomonella</i> on pear <i>E. Pasqualini, I. Espinha, S. Civolani, P. Medrzycki, E. Ladurner</i> | 399 |
| Control of the Grapevine Moth, <i>Lobesia botrana</i> (Den. et Schiff.) (Lepidoptera Tortricidae), by disorientation <i>Marco Tasin, Gianfranco Anfora, Gino Angeli, Mario Baldessari, Antonio de Cristofaro, Giacinto S. Germinara, Franco Rama, Silvia Vitagliano, Claudio Ioriatti</i> | 403 |
| Field evaluation of the sex pheromone of the Lackey moth <i>Malacosoma neustria</i> (L.) in two Italian regions <i>Giuseppe Rotundo, Omar V. Cau, Giacinto S. Germinara, Pietro Luciano, Antonio de Cristofaro</i> | 409 |
| Electrophysiological responses of two different species of apple gall midges (Diptera Cecidomyiidae) to host plant volatiles <i>Gianfranco Anfora, Claudio Ioriatti, Sergio Moser, Giacinto S. Germinara, Antonio de Cristofaro</i> | 413 |
| Effect of Madex® (granulovirus) on codling moth egg laying and larval damages on two apple varieties – Relationships with plant surface metabolites. <i>Nadia Lombarkia, Claudio Ioriatti, Sylvie Derridj</i> | 419 |
| Effects of the (E; Z)-2,4-ethyl decadienoate (DA2313) and synthetic pheromone blends to monitor <i>Cydia pomonella</i> adults <i>Michela Villa, Fabio Molinari, Edison Pasqualini, Isabel Espinha, Benoit Sauphanor</i> .. | 425 |
| Electrophysiological and olfactory responses of <i>Lobesia botrana</i> (Den. et Schiff.) (Lepidoptera Tortricidae) to odours of host plant <i>Silvia Vitagliano, Gianfranco Anfora, Marco Tasin, Giacinto S. Germinara, Claudio Ioriatti, Giuseppe Rotundo, Antonio De Cristofaro</i> | 429 |
| Allelochemical effect of the fruit trees, <i>Ziziphus spina christi</i> , on the searching behaviour of the parasitoid <i>Bracon hebetor</i> Say. (Hym., Braconidae) <i>Mourad Shonouda</i> | 437 |

ABSTRACTS

| | |
|--|-----|
| The state of the art <i>Carlo Malavolta</i> | 443 |
| Sustainable Winegrowing New Zealand: Technical developments and achievements <i>Sarah Gurnsey</i> | 444 |
| Genetic analysis of plum curculio, a fruit pest of export concern <i>Douglas G. Pfeiffer, Michelle McClanan, Zhang Xing Shirley Luckhart</i> | 445 |
| The ecological infrastructures of the farm to maintain and increase functional biodiversity <i>Ernst Boller, Jesús Avilla, Erich Jörg, Carlo Malavolta</i> | 446 |
| Peach extrafloral nectaries impact natural enemies and biological control of the oriental fruit moth <i>Clarissa R. Mathews, M. W. Brown</i> | 447 |
| Kaolin particle film as a pesticide in organic fruit production in a cool climate <i>Gunnhild Jaastad, Åse Marie Helgheim, Dag Røen, Arne Stensvand, Lars Olav Brandsæter</i> | 448 |
| Earwigs in stone fruit orchards <i>J. F. Mandrin</i> | 449 |
| The impact of Integrated Fruit Production on pest and disease management in New Zealand's apple industry <i>J.T.S. Walker, D.W.L Manktelow, M.B. Butcher</i> | 450 |
| Development in integrated pome fruit production in Portugal <i>J. Clemente</i> | 451 |

| | |
|--|-----|
| Information technology and the dissemination of fruit pest management information in Virginia, USA <i>Douglas G. Pfeiffer</i> | 452 |
| Predicting the risk of Brown rot in peaches after harvest <i>Jean-François Mandrin, Jean Lichou</i> | 453 |
| Susceptibility citrus leaf miner, <i>Phyllocnistis citrella</i> in Punjab, Pakistan, to different insecticides <i>Waqas Wakil, Mansoor-ul-Hasan, Faiza Bashir</i> | 454 |
| Preliminary study about the control of the leafhopper <i>Jacobiasca lybica</i> using a system with cover crop and mating disruption for <i>Lobesia botrana</i> <i>S. Garcia, C. Frescata</i> | 455 |
| Effect of groundcover management on apple orchard arthropods <i>V. Markó, K. Balázs, G. Jenser</i> | 456 |
| Use of Spray Adjuvants to enhance the performance of BOTRY-Zen® in grapes <i>S. Gurnsey, R. Gaskin, D. Manktelow, P. Elmer, F. Parry, P. Wood</i> | 457 |
| Results from a field study to monitor effects of Spinosad on the natural community in an apple orchard <i>Heidrun Vogt</i> | 458 |
| Effects of neem oil and NeemAzal® on the life-cycle of the vine moth, <i>Lobesia botrana</i> (Lepidoptera: Tortricidae) <i>Antonio Fenio, Massimo Cristofaro, Fortunata Minelli</i> | 459 |
| Prospects for developing eco-friendly biopesticides from natural plant species in Nigeria <i>A.C.Amadioha</i> | 460 |
| Production and Marketing of Organic Hazelnuts: the Case of “Tonda Gentile Romana” <i>Barbara Pancino and Silvio Franco</i> | 461 |
| Current Status of IFP Programs for Stone and Pome Fruits in Oregon, USA <i>Helmut Riedl, Steven P. Castagnoli, Lynn E. Long, and Michael K. Omeg</i> | 462 |
| Phenological model of pear psylla <i>Cacopsylla pyri</i> <i>L. Schaub, B. Graf, M. Hächler, A. Butturini</i> | 463 |
| Toxicity of five fungicides to predatory mites (Acari: Phytoseiidae) in an apple orchard in northern Portugal. <i>Raul Rodrigues, Joaquim Guerner-Moreira, Jorge Costa & Pedro Duarte</i> | 464 |
| The production of “green” and “yellow” lists in Integrated Production Guidelines <i>Ernst Boller, Jesús Avilla, Erich Jörg, Carlo Malavolta, Peter Esbjerg</i> | 465 |
| Adult <i>Cydia pomonella</i> L. responses to apple volatiles: sex, mating status and dosage effect <i>Daniel Casado, Jesús Avilla, César Gemenó, Magí Riba</i> | 466 |
| Laboratory and field tests to optimize the attraction of apple fruit moth (<i>Argyresthia conjugella</i> Zeller) to volatiles emitted from rowan (<i>Sorbus aucuparia</i> L.) <i>G.K. Knudsen, T. Hofsvang, S. Kobro, G. Jaastad, M. Bengtsson, P. Witzgall</i> | 467 |
| Occurrence of olfactory cells responding to pheromone components and plant volatile compounds in different species of Lepidoptera and Diptera: possible implications for semiochemical applications <i>A. de Cristofaro, G. Anfora, C. Ioriatti, G.S. Germinara, G. Rotundo</i> | 468 |
| Tactics to use mating disruption of codling moth in pear orchards of Oeste region in Portugal <i>J. Clemente</i> | 469 |
| Sex pheromone trapping of <i>Cossus cossus</i> (L.) and <i>Zeuzera pyrina</i> (L.) (Lepidoptera, Cossidae) <i>Paola Riolo, Sandro Nardi, Marzia Cristiana Rosi, Nunzio Isidoro, Stefano Maini</i> | 470 |
| Attractive plant volatiles to control the apple fruit moth (<i>Argyresthia conjugella</i> Zeller) <i>G. Jaastad, G.K. Knudsen, S. Kobro, P. Witzgall, M. Bengtsson, A.-C. Bäckman</i> | 471 |
| The sex pheromone of the apple leaf curling midge <i>Dasineura mali</i> <i>Jerry Cross</i> | 472 |
| Synthesis from germacrone of (±)-9-methylgermacrene, an active analogue of (S)-9- | |

| | |
|--|-----|
| methylgermacrene-B the sex pheromone of the sandfly, <i>Lutzomyia longipalpis</i> from Lapinha, Brazil <i>B. Krishnakumari, K. Sarita Raj, J.G.C. Hamilton</i> | 473 |
| Quantitative method for pheromone delivery in studies of sensory adaptation of moth antennae <i>R.M. Trimble, D.B. Marshall</i> | 474 |
| Olfactory sensitivity of <i>Bactrocera oleae</i> (Gmelin) (Diptera Tephritidae) adults to host- plant semiochemicals and kairomone identification by electrophysiological (EAG, SCR) and field studies <i>A. de Cristofaro, G. Rotundo, A. Belcari, G.S. Germinara</i> | 475 |
| Integrating mating disruption and new insecticides in Washington apple orchards, 2001-2003. <i>Jay F. Brunner, Elizabeth Beers, John Dunley, Vincent P. Jones</i> | 476 |
| Prevention of mating by auto confusion™: - New pest management technology using electrostatic powders <i>Philip Howse</i> | 477 |
| Effect of Plant Volatiles on the Growth of <i>Pythium aphanidermatum</i> , <i>Fusarium</i> <i>oxysporum</i> and <i>Botrytis cinerea</i> Strains Pathogenic on Kiwi and Grapevine <i>N. Varela, B. Alsanius, M. Bengtsson and P. Witzgall</i> | 479 |
| The Systematic and Efficient Use of Mating Disruption <i>Kinya Ogawa, Toshimi Kobayashi, Tatsuya Hojo</i> | 480 |
| Sampling methods in orchard trials: A comparison between beating and inventory sampling <i>Heidrun Vogt, Jutta Mütter</i> | 481 |
| A possibility to enhance natural enemies in apple orchards? – Reduced pesticide treatments in an apple orchard at East Malling, U.K <i>V. Markó, J. V. Cross, B. Keresztes, E. Kondorosy</i> | 482 |
| Effects of an azadirachtin-based compound on the host-parasitoid interactions between the Mediterranean fruit fly, <i>Ceratitis capitata</i> Wied. (Diptera: Tephritidae) and the braconid wasp <i>Opius concolor</i> Szepi. (Hymenoptera: Braconidae) <i>Giuseppe Carbone, Massimo Cristofaro, Vincenzo Di Ilio, Fortunata Minelli</i> | 483 |
| Current issues impacting on organic apple production in New Zealand <i>J.T.S. Walker, S. J. McArtney, P.N. Wood, J.N. Wunsche, J.W. Palmer</i> | 485 |